Preventing non-ventilator hospital-acquired pneumonia

Issue:

It's estimated that one in every 100 hospitalized patients will be affected by non-ventilator hospital-acquired pneumonia (NVHAP). While NVHAP is a significant patient safety and quality of care concern, it is not currently recognized as one of the National Database of Nursing Quality indicators for which hospitals are held accountable; nor is it one of the conditions that the Centers for Medicare & Medicaid Services (CMS) requires hospitals to report to the Centers for Disease Control & Prevention (CDC) National Healthcare Safety Network; and it is not integrated into the CMS current pay-for-reporting or performance programs.[1] As a result, this leaves NVHAP a health care-acquired condition without national tracking or accountability, and, most likely, is unaddressed by health care organizations.

A recent article in the journal Infection Control & Hospital Epidemiology (ICHE) detailed a call to action from national organizations, including The Joint Commission, to address NVHAP. The call to action includes launching a national health care conversation about NVHAP prevention and encouraging researchers to develop new strategies for NVHAP surveillance and prevention. This issue of Quick Safety focuses on the call's challenge to health care systems to implement and support NVHAP prevention, and to add NVHAP prevention measures to education for patients, health care professionals and students.[1]

Current NVHAP prevention strategies

Since the development of NVHAP requires a complex interaction of events that includes aspiration of microorganisms present in the oral cavity and a vulnerable host, most prevention measures target primary source control, and may include:[1,2]

- Maintaining regular oral care[1,2,3,4,5,6]
- Maintaining patient mobility[1,3,5,6,7]
- Elevating the head of the patient's bed[1,3,5,6]
- Reducing the use of acid-suppressing medications[1]
- Minimizing sedation[1,5]
- Performing dysphagia screening in high-risk patients[1]
- Using modified diets and feeding strategies for patients with abnormal swallowing[1,3,5]
- Following standardized processes to place and manage feeding tubes[1,5]

and labor intensive to apply. Despite these limitations and challe
medical centers can take to prevent NVHAP while improving the
the risk of sepsis, reducing health care costs, and saving lives.

1. Obtain buy-in from leadership and health care providers about the importance of NVHAP prevention.[1]
2. Overcome beliefs that NVHAP prevention strategies such as oral hygiene and mobility are optional tasks rather than standard-of-care interventions.[1]
3. Procure supplies necessary to implement effective interventions.[4]
4. Educate staff

Legal disclaimer: This material is meant as an information piece only; it is not a standard or a Sentinel Event Alert. The intent of Quick Safety is to raise awareness and to be helpful to Joint Commission-accredited organizations. The information in this publication is derived from actual events that occur in health care.

©2021 The Joint Commission, Division of Healthcare Improvement